

CP-Violating Yukawa Couplings in the Skyrme Model and the Neutron Electric Dipole Moment*

Harold A. Riggs and Howard J. Schnitzer

*Department of Physics, Brandeis University
Waltham, MA 02254*

November 1992

ABSTRACT

We argue that the large- N_c behaviour of the Yukawa couplings in the Skyrme model involves issues more subtle than the vanishing of linear fluctuations needed for classical stability of the skyrmion. The chiral fluctuations about the skyrmion must be quantized in order to reach a conclusion. An improved quantization procedure allows us to confront this question directly.

The pion-nucleon coupling constants $g_{\pi NN}$ (CP conserving) and $\bar{g}_{\pi NN}$ (CP violating) are calculated in the large- N_c , three-flavour Skyrme model by direct evaluation of the leading matrix elements appearing in the LSZ reduction formula. We find that $g_{\pi NN} \sim N_c^{\frac{3}{2}}$, but that, at most, $\bar{g}_{\pi NN} \sim m_\pi^2 N_c^{-\frac{1}{2}}$. These results show that the leading contribution to the neutron electric dipole moment in large- N_c Skyrme model is the direct one ($D_n \sim N_c m_\pi^2$), rather than the pion loop contribution.

*Supported in part by the DOE under grant DE-FG02-92ER40706

1. Introduction

Some time ago one of us considered chiral fluctuations about a rotating, three-flavour skyrmion¹

$$U(\mathbf{r}, t) = U_\phi(\mathbf{r}, t) A(t) U_0(\mathbf{r}) A^{-1}(t) U_\phi(\mathbf{r}, t) \quad (1.1)$$

where $U_0(\mathbf{r})$ is a static solution of the U(3) Skyrme model, $A(t)$ is an SU(3) matrix of collective coordinates, and the matrix field (summation convention throughout)

$$U_\phi = \exp\left\{\frac{i}{f_\pi} \phi_\alpha \lambda_\alpha\right\} \quad (1.2)$$

is built from the nonet of pseudoscalar meson fields ϕ_α ($\alpha = 0, 1, \dots, 8$) and the Gell-Mann matrices λ_a ($a = 1, \dots, 8$) with $\lambda_0 = \sqrt{2/3}\mathbf{I}$. Insertion of expression 1.1 into the Skyrme lagrangian, $\mathcal{L}_S$, yields the elegant result

$$\mathcal{L}_S(U(\mathbf{r}, t)) = \mathcal{L}_S(AU_0A^{-1}) + \mathcal{L}_\sigma(U_{2\phi}) + \mathcal{L}_I(AU_0A^{-1}; \phi) , \quad (1.3)$$

in which the right-hand side is isomorphic to the low-energy soft-meson and baryon effective lagrangian

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_B(\psi) + \mathcal{L}_\sigma(\pi) + \mathcal{L}_I(\psi; \pi) \quad (1.4)$$

which describes free baryons interacting with the mesons of the non-linear σ -model. Since 1.4 obeys all the soft-meson theorems of chiral symmetry, one expects that these theorems are obeyed by the skyrmion-meson system of 1.3 as well.

The Skyrme lagrangian gives a *formal* result for the Yukawa couplings of the skyrmion-meson interaction. However, to arrive at a conclusive result for the Yukawa couplings involves a number of subtleties which lead to the central issues of this paper. Given that 1.3 includes appropriate CP violating terms, one can compute both the CP violating pion-nucleon scalar coupling constant $\bar{g}_{\pi NN}$ and the CP conserving pseudoscalar coupling constant $g_{\pi NN}$ appearing in the interaction term of eq. 1.4

$$\mathcal{L}_I(\psi; \pi) = \boldsymbol{\pi} \cdot \bar{\psi} \boldsymbol{\tau} (i\gamma_5 g_{\pi NN} + \bar{g}_{\pi NN}) \psi . \quad (1.5)$$

Since the Skyrme model is formulated in terms of both chiral symmetry and the large N_c (number of colors) expansion, one expects that its predictions for $g_{\pi NN}$ and $\bar{g}_{\pi NN}$ are consistent with QCD in the same limits. Naïvely, the dependence of the couplings on the pion (nucleon) mass m_π (M_N) and the number of colors N_c is, from 1.3 and ref. 1,

$$g_{\pi NN} \sim N_c^{\frac{3}{2}} \quad \bar{g}_{\pi NN} \sim m_\pi^2 N_c^{\frac{1}{2}} \quad M_N \sim N_c . \quad (1.6)$$

Current algebra provides an ostensibly model-independent derivation of the leading contribution to the electric dipole moment of the neutron D_n in the chiral limit²

$$D_n = \frac{g_{\pi NN} \bar{g}_{\pi NN}}{4\pi^2 M_N} \ln\left(\frac{M_N}{m_\pi}\right) . \quad (1.7)$$

If eq. 1.6 were correct, the leading contribution in the combined chiral and large N_c limit (and so dominant prediction of the Skyrme model in these limits) would be $D_n \sim N_c m_\pi^2 \ln m_\pi$.

Recently significant progress in resolving this issue has been made in calculating $g_{\pi NN}$ (in the two flavour Skyrme model) by the authors of ref. 3. It is the purpose of this paper to calculate the CP conserving and CP violating π -N couplings $g_{\pi NN}$, and $\bar{g}_{\pi NN}$ by the methods of ref. 3, generalized to the three-flavour Skyrme model. We will show that although the pseudoscalar coupling $g_{\pi NN}$ indeed has the leading behaviour of eq. 1.6, the CP violating coupling $\bar{g}_{\pi NN}$ is down by a factor of at least N_c^{-1} from that of eq. 1.6. Therefore, the pion contribution to D_n is at most

$$D_n \sim m_\pi^2 \ln m_\pi \quad (1.8)$$

and is therefore subdominant to the direct CP violating behaviour $D_n \sim m_\pi^2 N_c$ computed in ref. 4. (For a review see ref. 5.)

It is important to understand why there is an issue with regard to the Skyrme model Yukawa couplings. These couplings embody the linear fluctuations about the Skyrme classical solution. For constant collective coordinates ($A = \text{constant}$), this linear coupling must vanish for the classical solution to be stable under linear fluctuations. Thus the *classical*, linear meson-skyrmion interactions behave as

$$\mathcal{L} \sim \phi (\dot{A} \text{ terms}) \quad (1.9)$$

if the Skyrme equations of motion are imposed. The time derivatives in eq. 1.9 imply a suppression of at least a factor of N_c^{-1} compared to the behaviour given in 1.6. If the chiral fluctuations are only considered classically, this would be the end of the story. However, the classical argument cannot be trusted since the chiral fluctuations must be quantized in order to properly account for the mesons as quantum fields.^{6, 7} This is the origin of a number of complications.

The canonical momenta of the rotating skyrmion and associated mesons are not independent, and the separation into components transverse to the zero-mode solutions is “gauge dependent.” Correctly calculating an S -matrix element—which must be gauge invariant—such as the π -N scattering amplitude is a complicated task.⁸ The advance made by the authors of ref. 3 is to calculate the Born terms of π -N scattering in a manifestly gauge-invariant manner in the (two-flavour) Skyrme model. They obtain the same result as that given by the gauge-dependent separation of skyrmion and meson momenta,⁸ confirming that the Yukawa coupling $g_{\pi NN}$ has the large N_c behaviour of eq. 1.6. This suggests that the large- N_c analysis of CP conserving π -N scattering in the Skyrme model is consistent with the implications of chiral symmetry.

The gauge-invariant calculation presented in ref. 3 is considerably simpler than the calculations in a gauge that separates the skyrmion and meson momenta. As a result, application of this method to the calculation of the $\pi - N$ couplings in the three-flavour Skyrme model yields a straightforward estimate of the large- N_c behaviour of $g_{\pi NN}$ and $\bar{g}_{\pi NN}$.

2. The CP Violating Skyrmion

In order to obtain strong CP violation, one must consider a U(3) non-linear σ -model so as to include (i) an effective interaction representing the U(1) anomaly and (ii) a CP violating mass-matrix. The appropriate effective lagrangian in the large N_c limit is then⁹

$$\mathcal{L}(U) = \mathcal{L}_S(U) + \frac{f_\pi^2}{16} \text{Tr}(MU + M^\dagger U^\dagger - M - M^\dagger) + \frac{af_\pi^2}{64N_c} (\text{Tr}[\ln U - \ln U^\dagger])^2 + \mathcal{L}_{WZ} \quad (2.1)$$

where $\mathcal{L}_{WZ}$ is the Wess-Zumino 5-form, U is a U(3) matrix field, a depends on the physical meson masses,

$$\mathcal{L}_S(U) = \frac{f_\pi^2}{16} \text{Tr}[(\partial_\mu U)(\partial_\mu U^\dagger)] + \frac{1}{32} e^{-2} \text{Tr}\{[(\partial_\mu U)U^\dagger, (\partial_\nu U)U^\dagger]^2\} \quad (2.2)$$

and

$$M = \text{diag}[M_1 \exp(i\frac{\beta_1}{M_1}), M_2 \exp(i\frac{\beta_2}{M_2}), M_3 \exp(i\epsilon\frac{\beta_3}{M_3})] . \quad (2.3)$$

In order to prevent strong CP violation from creating Goldstone bosons from the vacuum we require^{2, 4}

$$\begin{aligned} \beta_1 = \beta_2 = \beta_3 &\equiv \beta \\ \epsilon &= \frac{a + M_3}{a + M_1} \end{aligned} \quad (2.4)$$

up to terms of $\mathcal{O}(\beta^2)$. It is convenient to define $\theta = \arg \det M$, the effective CP angle, and to set

$$\theta = \lambda_\theta \left(\frac{1}{M_1} + \frac{1}{M_2} + \frac{\epsilon}{M_3} \right) \quad (2.5)$$

so that $\lambda_\theta = \beta$.

The classical SU(3) Skyrme ansatz, neglecting CP violation, is^{10, 11}

$$U_0(\mathbf{r}) = \begin{pmatrix} \cos F + i\boldsymbol{\tau} \cdot \mathbf{r} \sin F & 0 \\ 0 & 0 & 1 \end{pmatrix} = \frac{\sqrt{2}}{3} (\cos F - 1) \lambda_8 + \frac{1}{3} (2 \cos F + 1) I + i\boldsymbol{\tau} \cdot \mathbf{r} \sin F \quad (2.6)$$

where $F = F(r)$. As emphasized by the authors of ref. 4 the CP violating term in $\mathcal{L}$ reacts back on the skyrmion. We adopt their generalization of ansatz 2.6 and consider

$$U_c(\mathbf{r}) = \exp\left(\frac{2i}{f_\pi} \left[\sqrt{\frac{2}{3}} (f_0(r) - \delta) I + \lambda_8 f_8(r) \right] \right) U_0(\mathbf{r}) \quad (2.7)$$

where

$$\delta = \frac{\sqrt{6}}{4} \frac{\lambda_\theta f_\pi}{a + m_\pi^2}$$

is the further vacuum shift required in order to prevent the creation of Goldstone bosons from the vacuum.⁴ The ansatz in eq. 2.7 remains spherically symmetric under combined isospin and spatial rotations. Neglecting isospin violation one can take

$$M_1 = M_2 = m_\pi^2 \quad M_3 = 2m_K^2 - m_\pi^2 \quad 3a/N_c = m_\eta^2 + m_{\eta'}^2 - 2m_K^2 \quad (2.8)$$

so that $\lambda_\theta \sim \frac{1}{2}m_\pi^2\theta$. Since it does not affect any issues of principle, we neglect η - η' mixing and corrections of order $\mathcal{O}(M_1/a)$ in order to simplify the discussion. Taking these effects into account would only produce small *numerical* corrections of our results. The CP violating (CPV) part of the lagrangian, which contains contributions from both the mass and anomaly terms,⁴ provides a source for $f_0(r)$ and $f_8(r)$:

$$\delta\mathcal{L}_{CPV} = \frac{\lambda_\theta f_\pi}{2\sqrt{3}}(1 - \cos F)(f_8 + \sqrt{2}f_0) + \mathcal{O}\left(\frac{M_1}{a}\right). \quad (2.9)$$

The function $F(r)$ is unchanged to order $\mathcal{O}(\theta)$ by the presence of this perturbation.

Neglecting terms of order $\mathcal{O}(m_\pi^2/m_{\eta'}^2)$ one can use this source to solve for f_0 and f_8 in terms of $F(r)$. The result is⁴

$$\begin{aligned} f_8(r) &= \frac{\lambda_\theta f_\pi}{2\sqrt{3}} \int \frac{d^3r'}{4\pi} \frac{e^{-m|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|} [1 - \cos F(r')] \\ f_0(r) &= \frac{\sqrt{2}\lambda_\theta f_\pi}{2\sqrt{3}} \int \frac{d^3r'}{4\pi} \frac{e^{-m'|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|} [1 - \cos F(r')] \end{aligned} \quad (2.10)$$

where we have neglected η - η' mixing, and have written $m = m_\eta$ and $m' = m_{\eta'}$. The angular integration in eqs. 2.10 gives

$$\begin{aligned} f_8(r) &= \frac{\lambda_\theta f_\pi}{2\sqrt{3}m} \int_0^\infty dr' G_m(r, r') [1 - \cos F(r')] \\ f_0(r) &= \frac{\sqrt{2}\lambda_\theta f_\pi}{2\sqrt{3}m'} \int_0^\infty dr' G_{m'}(r, r') [1 - \cos F(r')] \end{aligned} \quad (2.11)$$

where

$$G_m(r, r') = \begin{cases} \left(\frac{r'}{r}\right) e^{-mr'} \sinh(mr) & r < r' \\ \left(\frac{r'}{r}\right) e^{-mr} \sinh(mr') & r > r' \end{cases} \quad (2.12)$$

The contributions of $f_8(r)$ and $f_0(r)$ will be important in what follows.

3. Meson-Nucleon Scattering in the Skyrme Model

Here we generalize the analysis of ref. 3 to the CP-violating, three-flavour model described in the previous section. Denote the meson field (its quantum nature signaled by the absence of the subscript c) by

$$U(\mathbf{r}, t) = \lambda_\alpha U_\alpha \quad \alpha = 0, 1, \dots, 8. \quad (3.1)$$

It is convenient to write

$$U_\alpha = s_\alpha + ip_\alpha \quad (3.2)$$

where s_α , and p_α are *real* fields. The non-linear σ model constraint that U is a $U(3)$ field,

$$U^\dagger U = I, \quad (3.3)$$

gives nine constraint equations. Therefore we can regard the s_α fields as dependent variables expressible in terms of the dynamical fields p_α . (These nine independent real fields embody the pseudoscalar nonet.)

Substituting eq. 3.1 into the lagrangian (eq. 2.1) we obtain (exhibiting all time derivatives explicitly)

$$\mathcal{L} = \frac{1}{2} \dot{p}_\alpha K_{\alpha\beta} \dot{p}_\beta + W_\alpha \dot{p}_\alpha - \mathcal{V}(p, \nabla p) \quad (3.4)$$

where $\alpha, \beta = 0, 1, \dots, 8$ and W_α is obtained from the Wess-Zumino term. (This is the analogue of eq. 3.3 of ref. 3.) In the weak field limit

$$K_{\alpha\beta} = \frac{f_\pi^2}{4} \delta_{\alpha\beta} + \dots \quad (3.5)$$

Beyond this, the detailed forms of $K_{\alpha\beta}$, W_α , and $\mathcal{V}(p, \nabla p)$ will not be needed. The canonical momenta are

$$\Pi_\alpha = \frac{\delta L}{\delta \dot{p}_\alpha} = K_{\alpha\beta} \dot{p}_\beta + W_\alpha \quad (3.6)$$

so that

$$\dot{p}_\alpha = (K^{-1})_{\alpha\beta} (\Pi_\beta - W_\beta) . \quad (3.7)$$

Consequently, the Hamiltonian is, up to terms of $\mathcal{O}(\hbar^2)$ since we have ignored the problem of operator ordering,

$$\mathcal{H} = \frac{1}{2} \int d^3x (\Pi - W)_\alpha (K^{-1})_{\alpha\beta} (\Pi - W)_\beta + \mathcal{V}(p, \nabla p) \quad (3.8)$$

and the canonical commutation relations are

$$[p_\alpha(x), \Pi_\beta(y)] \delta(x^0 - y^0) = i \delta_{\alpha\beta} \delta^{(4)}(x - y) . \quad (3.9)$$

Due to the derivative couplings $K_{\alpha\beta} \neq \frac{f_\pi^2}{4} \delta_{\alpha\beta}$ and $\mathcal{H}_{\text{int}} \neq -\mathcal{L}_{\text{int}}$, as is evident from eq. 3.4-3.9. This is the essential reason that the classical argument for the absence of Yukawa couplings described in the introduction fails.

The skyrmion soliton is the *static* but *rotated* solution (*i.e.*, $U^s = AU_c A^{-1}$) to the equations of motion obtained from eq. 3.4. (The superscript s will denote the static, classical part of a quantity in the following.) In terms of the field of eq. 2.7, this solution is given by

$$p_\alpha^s = \frac{1}{2} \text{Im Tr}(\lambda_\alpha A U_c A^{-1}) \quad (3.10)$$

where $A \in \text{SU}(3)$ rotates the ansatz U_c to an arbitrary direction in $\text{SU}(3)$ space and for the static solution we neglect any time dependence of A .

In order to obtain the S-matrix for the meson-nucleon elastic scattering, we will analyze the LSZ reduction formula. To carry this out we must identify the interpolating field Φ_α . In fact, it is proportional to the quantum field p_α in eq. 3.4 (not the classical p_α^s). Since $\lim_{r \rightarrow \infty} U_c(r) = \text{I}$, we can pick out the proportionality constant from the non-linear meson fields which appear in eqs. 1.2 & 1.3 by using the fact that $\lim_{r \rightarrow \infty} U(\mathbf{r}, t) = U_{2\phi}$:

$$U_{2\phi} \sim 1 + \frac{2i}{f_\pi} \lambda \cdot \phi + \mathcal{O}\left(\frac{\phi^2}{f_\pi^2}\right) \quad (3.11)$$

so that

$$\Phi_\alpha = \frac{f_\pi}{2} p_\alpha. \quad (3.12)$$

Therefore

$$\Phi_\alpha = \frac{f_\pi}{4} \text{ImTr}[\lambda_\alpha U(\mathbf{r}, t)] \quad (3.13)$$

where $U(\mathbf{r}, t)$ is the complete field $U_\phi A U_c A^{-1} U_\phi$ that enters $\mathcal{L}(U)$.

The reduction formula for our problem is given in terms of the source term $\mathcal{J}(x)$, Φ and $\dot{\Phi}$, as follows

$$\begin{aligned} S_{fi} &= \delta_{fi} + i^2 \int d^3\mathbf{r}' dt' \int d^3\mathbf{r} dt f_{k'}^*(x') f_k(x) \\ &\times \left\langle B'(\mathbf{p}') | T(\mathcal{J}_\beta(x') \mathcal{J}_\alpha(x)) + \delta(t' - t) [\dot{\Phi}_\beta(x'), \mathcal{J}_\alpha(x)] - i\omega_{k'} \delta(t' - t) [\Phi_\beta(x'), \mathcal{J}_\alpha(x)] | B(\mathbf{p}) \right\rangle \end{aligned} \quad (3.14)$$

with $f_{\mathbf{k}}(x) = e^{-ik \cdot x} / \sqrt{(2\pi)^3 2\omega_k}$, where $k \cdot x = \omega_k t - \mathbf{k} \cdot \mathbf{r}$. The source term of the meson fields is

$$\mathcal{J}_\alpha(x) = -[H, [H, \Phi_\alpha(x)]] + (-\nabla^2 + m_\pi^2) \Phi_\alpha(x) \quad (3.15)$$

with the commutator computed using eq. 3.8, 3.9, and 3.12. The result is

$$\begin{aligned} [H, [H, p_\alpha(y)]] &= (K^{-1})_{\alpha\beta} \frac{\delta \mathcal{V}(\mathbf{p}, \nabla p)}{\delta p_\beta} \\ &+ \frac{1}{2} (\Pi - W)_\gamma (K_{\alpha\beta}^{-1} K_{\gamma\delta, \beta}^{-1} - 2K_{\gamma\beta}^{-1} K_{\alpha\delta, \beta}^{-1}) (\Pi - W)_\delta \\ &+ (\Pi - W)_\gamma K_{\alpha\beta}^{-1} K_{\gamma\delta}^{-1} (W_{\beta, \delta} - W_{\delta, \beta}) \end{aligned} \quad (3.16)$$

where for any quantity T and any index α , the notation $T_{,\alpha}$ denotes $\delta T / \delta p_\alpha$. We have not kept track of operator orderings, so that eq. 3.16 is accurate only up to terms of $\mathcal{O}(\hbar^2)$. We again see that the presence of derivative couplings in $\mathcal{L}$ and $\mathcal{H}$ produces a different result in eq. 3.16 than that expected from classical considerations.

Similarly, using eq. 3.15 & 3.16, we find that

$$\begin{aligned}
[\dot{p}_\beta(x), \mathcal{J}_\alpha(y)]\delta(x^0 - y^0) &= -i(K^{-1})_{\beta\gamma} \frac{\delta \mathcal{J}_\alpha(y)}{\delta p_\gamma(x)} \delta(x^0 - y^0) \\
&+ [(K^{-1})_{\beta\gamma}, \mathcal{J}_\alpha(y)](\Pi - W)_\gamma \delta(x^0 - y^0) \\
&- (K^{-1})_{\beta\gamma} [W_\gamma(x), \mathcal{J}_\alpha(y)] \delta(x^0 - y^0)
\end{aligned} \tag{3.17}$$

with

$$[\dot{\Phi}_\beta(x), \mathcal{J}_\alpha(y)]\delta(x^0 - y^0) = \frac{f_\pi}{2} [\dot{p}_\beta(x), \mathcal{J}_\alpha(y)]\delta(x^0 - y^0) \tag{3.18}$$

We also need

$$[\Phi_\beta(x), \mathcal{J}_\alpha(y)]\delta(x^0 - y^0) = i \frac{f_\pi}{2} \frac{\delta \mathcal{J}_\alpha(y)}{\delta \Pi_\beta(x)} \delta(x^0 - y^0) . \tag{3.19}$$

The above equations are again accurate only to order $\hbar$.

In order to evaluate the Born terms from eq. 3.14 we need certain one-baryon matrix elements, which will be analyzed in terms of the $1/N_c$ expansion.

The *crucial* point is that in the matrix elements of operators taken between single-baryon states, one may replace the field operators with the classical fields evaluated with the skyrmion solution (3.10), *i.e.*, $p_\alpha(x) \rightarrow p_\alpha^s(x)$. This idea goes all the way back to reference 12. The reason is that the fluctuating parts vanish since the “in” and “out” meson states form a Fock space on the one-baryon subspace. As emphasized in ref. 3 it is not necessary to explicitly separate Φ_α into static and fluctuating parts, since the one-baryon meson matrix elements will automatically pick out the static solution.

In order to arrange the one-baryon matrix elements in powers of $1/N_c$, we need to order the various classical field quantities in terms of powers of N_c . By the classical equations of motion

$$\frac{\delta \mathcal{V}(p^s, \nabla p^s)}{\delta p_\alpha^s} = 0 \tag{3.20}$$

Note that the ordering in $1/N_c$ is the *same* for the CP conserving (CPC) and CP violating (CPV) terms in p_α^s , to leading order in CP violation, as can be seen from the following. From eq. 3.10, the CP conserving term in p_α^s is

$$(p_\alpha^s)_{\text{CPC}} = \frac{1}{2} \text{tr}(\lambda_\alpha A \boldsymbol{\tau} A^{-1}) \cdot \mathbf{r} \sin F(r) \tag{3.21}$$

so that

$$(p_\alpha^s)_{\text{CPC}} \sim \mathcal{O}(1) \tag{3.22}$$

in the $1/N_c$ expansion. Then,

$$\begin{aligned}
p_\alpha^s &= (p_\alpha^s)_{\text{CPC}} + (p_\alpha^s)_{\text{CPV}} \\
&\sim \mathcal{O}(1)[1 + \lambda_\theta/m_\eta^2] \\
&\sim \mathcal{O}(1)[1 + (\frac{m_\pi^2}{m_\eta^2})\theta]
\end{aligned} \tag{3.23}$$

where we have used eqs. 2.7 and 2.11. Therefore, the $1/N_c$ estimates that follow hold in both the CP violating and conserving cases.

The estimates in eqs. 3.22 and 3.23, and eq. 3.9 imply that

$$(\Pi_s) \sim \mathcal{O}(1) \tag{3.24}$$

Since

$$(K_s)_{\alpha\beta} \sim \mathcal{O}(f_\pi^2) \sim \mathcal{O}(N_c) \tag{3.25}$$

one also has

$$\begin{aligned}
\dot{p}_\alpha^s &\sim \mathcal{O}(N_c^{-1}) \\
(K^{-1})_{\alpha\beta}^s &\sim \mathcal{O}(N_c^{-1}) \\
(W^s)_\alpha &\sim \mathcal{O}(1)
\end{aligned} \tag{3.26}$$

for these classical quantities. With these ingredients we can now make $1/N_c$ estimates of the one-baryon matrix elements.

For the first term in eq. 3.14 we need

$$\begin{aligned}
\langle B | \mathcal{J}_\alpha(x) | B' \rangle &= \langle B | (-\nabla^2 + m_\pi^2) \Phi_\alpha + \partial_t^2 \Phi_\alpha | B' \rangle \\
&= \frac{f_\pi}{2} \{ \langle B | (-\nabla^2 + m_\pi^2) (p_s)_\alpha | B' \rangle + \mathcal{O}(1/N_c^2) \}
\end{aligned} \tag{3.27}$$

which holds because all the terms in eq. 3.16, evaluated with the classical solution, contribute at most to the order $\mathcal{O}(1/N_c^2)$ correction.

Similarly, from eq. 3.19

$$\begin{aligned}
\langle B | [\Phi_\beta(x), \mathcal{J}_\alpha(y)] \delta(x^0 - y^0) | B' \rangle &= i \frac{f_\pi}{2} \langle B | \left[\frac{\delta \mathcal{J}_\alpha(y)}{\delta \Pi_\beta(x)} \right]_s | B' \rangle \delta(x^0 - y^0) \\
&\sim \mathcal{O}(f_\pi^{-3}) \sim \mathcal{O}(N_c^{-\frac{3}{2}})
\end{aligned} \tag{3.28}$$

and from eq. 3.18

$$\begin{aligned}
\langle B | [\dot{\Phi}_\beta(x), \mathcal{J}_\alpha(y)] \delta(x^0 - y^0) | B' \rangle &= \frac{f_\pi}{2} \{ -i \langle B | (K_s^{-1})_{\beta\gamma} \left[\frac{\delta \mathcal{J}_\alpha(y)}{\delta p_\gamma(x)} \right]_s | B' \rangle \delta(x^0 - y^0) + \mathcal{O}(1/N_c^2) \} \\
&\sim \mathcal{O}(1)
\end{aligned} \tag{3.29}$$

Therefore only eq. 3.27 and 3.29 contribute to the Born terms in the large N_c limit. In addition, only equation 3.27 appears in the pole term, and so gives both the CP conserving and CP violating Yukawa couplings, while eq. 3.29 contributes to a non-pole contact term.

4. The π -N Coupling Constants

The classical, CP-violating solution is, from eqs. 2.7, 3.12, and 3.13,

$$(\Phi_\alpha^s)_{\text{CPV}} = \frac{1}{30\sqrt{3}} \left\{ [\sqrt{2}(f_0(r) - \delta) + f_8(r)](\cos F(r) - 1) + 4f_8(r) \right\} \text{Tr}(\lambda_\alpha A \lambda_8 A^{-1}) \quad (4.1)$$

Therefore, to leading order in $1/N_c$,

$$\begin{aligned} \bar{g}_{\pi NN}(\mathbf{k}) &= \int \frac{d^3r}{(2\pi)^3} e^{-i\mathbf{k}\cdot\mathbf{r}} \langle B | (\mathcal{J}_\alpha)_{\text{CPV}} | B' \rangle \\ &\simeq \int \frac{d^3r}{(2\pi)^3} e^{-i\mathbf{k}\cdot\mathbf{r}} \langle B | (-\nabla^2 + m_\pi^2) (\Phi_\alpha^s)_{\text{CPV}} | B' \rangle \end{aligned} \quad (4.2)$$

If the physical (mass-shell) coupling

$$\bar{g}_{\pi NN} = \bar{g}_{\pi NN}(\mathbf{k})|_{\mathbf{k}^2 = -m_\pi^2} \quad (4.3)$$

is to have a contribution from eq. 4.2, then $(\Phi_\alpha^s)_{\text{CPV}}$ must have a pion tail (*i.e.* it must behave as $\exp(-m_\pi r)/r$ as $r \rightarrow \infty$). Since

$$F(r) \longrightarrow C \frac{e^{-m_\pi r}}{r} \quad \text{as } r \rightarrow \infty \quad (4.4)$$

it is clear that the $\cos F(r)$ factor does not have a pion tail. Further, eq. 2.10 shows that $f_8(r)$ and $f_0(r)$ have (momentum space) poles at m_η^2 and $m_{\eta'}^2$, respectively, rather than at m_π^2 , so that the required pion tail is entirely absent from eq. 4.1. By contrast, the CP conserving coupling is given by (the analysis is essentially the same as that already discussed in ref. 3)

$$g_{\pi NN}(\mathbf{k}) = \frac{2}{3} M_N f_\pi \frac{\mathbf{k}^2 + m_\pi^2}{|\mathbf{k}|} \int d^3r j_1(kr) \sin F(r) \quad (4.5)$$

where $j_1(y)$ is the spherical Bessel function. Since $\sin F(r)$ ($\sim F(r)$ as $r \rightarrow \infty$) *does* have the required pion tail, $g_{\pi NN}$ is non-vanishing on-shell.

Therefore we conclude that $\bar{g}_{\pi NN}$ is suppressed relative to eq. 1.6 by at least a factor of N_c^{-1} , while the behaviour of $g_{\pi NN}$ is correctly given by 1.6.[†] It is important

[†]Although the order $\mathcal{O}(N_c^{\frac{1}{2}})$ contribution to $\bar{g}_{\pi NN}$ does not vanish off-shell, it is suppressed by the factor m_π^2/m_η^2 , as is evident from eq. 3.23.

to note that since $g_{\pi NN}$ is not suppressed by factors of N_c^{-1} , the stability of the linear fluctuations about the classical solution cannot be the reason for the suppression of $\bar{g}_{\pi NN}$. The correct explanation rests with the asymptotic behaviour of the skyrmion profile function eq. 2.6, where one should compare the coefficient of λ_8 with that of τ .

5. Conclusions

Our calculation shows that the Yukawa couplings in the Skyrme model involve issues more subtle than the vanishing of linear fluctuations about the classical solution. The CP conserving coupling $g_{\pi NN}$ is not suppressed, while the CP violating coupling is. Although this supports the final result of ref. 4, we find that neglect of the pion contribution to D_n in the large N_c limit requires more analysis than indicated there before one can be assured that the direct contribution to D_n dominates. The behaviour of the CP violating coupling constant $\bar{g}_{\pi NN}$ is also interesting for other applications.¹³

The primary value of the Skyrme model is as a phenomenological approximation to the low-energy effective lagrangian of QCD; therefore, one hopes that it substantially embodies the physics of chiral symmetry and the large- N_c properties of QCD. In fact, we have found that simple QCD estimates based on naïve quark-line counting (*i.e.* estimates similar to those of ref. 14 but using the $\theta g^2 F \tilde{F}$ interaction) agree with our Skyrme model results. In regard to this, it would have been peculiar if the Skyrme model, which does give the correct large N_c behaviour for the multi-meson skyrmion, did not, in the end, give the correct Yukawa couplings as well.

References

- [1] H. J. Schnitzer, *Phys. Lett.* **B139** (1984) 217.
- [2] V. Baluni, *Phys. Rev.* **D19** (1979) 2227; R. Crewther, P. Di Vecchia, G. Veneziano, and E. Witten, *Phys. Lett.* **B88** (1979) 123; *errata* **B91** (1980) 487.
- [3] A. Hayashi, S. Saito, and M. Uehara, *Phys. Rev.* **D11** (1992) 4856.
- [4] L. Dixon, A. Larignau, Y. Nir, and B. Warr, *Phys. Lett.* **B253** (1991) 459.
- [5] Y. Nir, “CP Violation,” SLAC preprint, SLAC-PUB-5874, September 1992.
- [6] H. J. Schnitzer, *Nucl. Phys.* **B261** (1985) 546.
- [7] H. J. Schnitzer, *Phys. Lett.* **B253** (1991) 465.

- [8] A. Hayashi, S. Saito, and M. Uehara, *Phys. Lett.* **B246** (1990) 15; *Phys. Rev.* **D43** (1991) 1520; M. Uehara, A. Hayashi, and S. Saito, *Nucl. Phys.* **A534** (1991) 680.
- [9] C. Rosenzweig, J. Schechter, and C. Trahern, *Phys. Rev.* **D21** (1980) 3388; E. Witten, *Ann. Phys.* **128** (1980) 363; P. Di Vecchia, F. Nicodemi, R. Pettorino, and G. Veneziano, *Nucl. Phys.* **B181** (1981) 318.
- [10] G. Adkins, C. Nappi, and E. Witten, *Nucl. Phys.* **B228** (1983) 552; G. Adkins and C. Nappi, *Nucl. Phys.* **B233** (1984) 109.
- [11] E. Guadagnini, *Nucl. Phys.* **B236** (1984) 35.
- [12] J. Goldstone and R. Jackiw, *Phys. Rev.* **D11** (1975) 2943. J.-L. Gervais, A. Jevicki, and B. Sakita, *Phys. Rev.* **D12** (1975) 1038. E. Tomboulis, *Phys. Rev.* **D12** (1975) 1678.
- [13] W. Fischler, S. Paban, and S. Thomas, *Phys. Lett.* **B289** (1992) 373.
- [14] H. Mütter, C. Engelbrecht, and G. Brown, *Nucl. Phys.* **A462** (1987) 701.